

## Exercise 2.8

### Question # 1

Operation  $\oplus$  performed on the two-member set  $G = \{0,1\}$  is shown in the adjoining table. Answers the questions:

- Name the identity element if it exists?
- What is the inverse of 1?
- Is the set  $G$ , under the given operation a group?  
Abelian and non-abelian?

$\oplus$	0	1
0	1	1
1	1	0

### Solutions

- From the given table we have  
 $0+0=0$  and  $0+1=1$   
This show that 0 is the identity element.
- Since  $1+1=0$  (identity element) so the inverse of 1 is 1.
- It is clear from table that element of the given set satisfy closure law, associative law, identity law and inverse law thus given set is group under  $\oplus$ .  
Also it satisfies commutative law so it is an abelian group.

### Question # 2

The operation  $\oplus$  as performed on the set  $\{0,1,2,3\}$  is shown in the adjoining table, shown that the set is an Abelian group?

### Solution

Suppose  $G = \{0,1,2,3\}$

- The given table show that each element of the table is a member of  $G$  thus closure law holds.
- $\oplus$  is associative in  $G$ .
- Table show that 0 is identity element w.r.t.  $\oplus$ .
- Since  $0+0=0$ ,  $1+3=0$ ,  $2+2=0$ ,  $3+1=0$   
 $\Rightarrow 0^{-1}=0$ ,  $1^{-1}=3$ ,  $2^{-1}=2$ ,  $3^{-1}=1$

$\oplus$	0	1	2	3
0	0	1	2	3
1	1	2	3	0
2	2	3	0	1
3	3	0	1	2

- As the table is symmetric w.r.t. to the principal diagonal. Hence commutative law holds.

### Question # 3

For each of the following sets, determine whether or not the set forms a group with respect to the indicated operation. From above table solve these (i-v) options.

### Solution

- As  $0 \in \mathbb{Q}$ , multiplicative inverse of 0 is not in set  $\mathbb{Q}$ . Therefore the set of rational number is not a group w.r.t to " $\cdot$ ".
- Closure property holds in  $\mathbb{Q}$  under  $+$  because sum of two rational number is also rational.
  - Associative property holds in  $\mathbb{Q}$  under addition.
  - $0 \in \mathbb{Q}$  is an identity element.

d- If  $a \in \mathbb{Q}$  then additive inverse  $-a \in \mathbb{Q}$  such that  $a + (-a) = (-a) + a = 0$ .

Therefore the set of rational number is group under addition.

(iii) a- Since for  $a, b \in \mathbb{Q}^+$ ,  $ab \in \mathbb{Q}^+$  thus closure law holds.

b- For  $a, b, c \in \mathbb{Q}$ ,  $a(bc) = (ab)c$  thus associative law holds.

c- Since  $1 \in \mathbb{Q}^+$  such that for  $a \in \mathbb{Q}^+$ ,  $a \times 1 = 1 \times a = a$ . Hence 1 is the identity element.

d- For  $a \in \mathbb{Q}^+$ ,  $\frac{1}{a} \in \mathbb{Q}^+$  such that  $a \times \frac{1}{a} = \frac{1}{a} \times a = 1$ . Thus inverse of  $a$  is  $\frac{1}{a}$ .

Hence  $\mathbb{Q}^+$  is group under addition.

(iv) Since  $\mathbb{Z} = \{0, \pm 1, \pm 2, \pm 3, \dots\}$

a- Since sum of integers is an integer therefore for  $a, b \in \mathbb{Z}$ ,  $a + b \in \mathbb{Z}$ .

b- Since  $a + (b + c) = (a + b) + c$  thus associative law holds in  $\mathbb{Z}$ .

c- Since  $0 \in \mathbb{Z}$  such that for  $a \in \mathbb{Z}$ ,  $a + 0 = 0 + a = a$ . Thus 0 an identity element.

d- For  $a \in \mathbb{Z}$ ,  $-a \in \mathbb{Z}$  such that  $a + (-a) = (-a) + a = 0$ . Thus inverse of  $a$  is  $-a$ .

(v) Since  $\mathbb{Z} = \{0, \pm 1, \pm 2, \pm 3, \dots\}$

For any  $a \in \mathbb{Z}$  the multiplicative inverse of  $a$  is  $\frac{1}{a} \notin \mathbb{Z}$ . Hence  $\mathbb{Z}$  is not a group under multiplication.

#### Question # 4

Show that the adjoining table represents the sums of the elements of the set  $\{E, O\}$ .

What is the identity element of this set? Show that this set is abelian group.

#### Solution

As  $E + E = E$ ,  $E + O = O$ ,  $O + O = E$

Thus the table represents the sums of the elements of set  $\{E, O\}$ .

The identity element of the set is E because

$$E + E = E + E = E \quad \& \quad E + O = O + E = E.$$

$\oplus$	E	O
E	E	O
O	O	E

i) From the table each element belong to the set  $\{E, O\}$ .

Hence closure law is satisfied.

ii)  $\oplus$  is associative in  $\{E, O\}$

iii) E is the identity element of w.r.t to  $\oplus$

iv) As  $O + O = E$  and  $E + E = E$ , thus inverse of O is O and inverse of E is E.

v) As the table is symmetric about the principle diagonal therefore  $\oplus$  is commutative.

Hence  $\{E, O\}$  is abelian group under  $\oplus$ .

#### Question # 5

Show that the set  $\{1, \omega, \omega^2\}$ , when  $\omega^3 = 1$  is an abelian group w.r.t. ordinary multiplication.

#### Solution

Suppose  $G = \{1, \omega, \omega^2\}$

$\otimes$	1	$\omega$	$\omega^2$
1	1	$\omega$	$\omega^2$
$\omega$	$\omega$	$\omega^2$	1
$\omega^2$	$\omega^2$	1	$\omega$

- i) A table show that all the entries belong to  $G$ .
- ii) Associative law holds in  $G$  w.r.t. multiplication.  
 e.g.  $1 \times (\omega \times \omega^2) = 1 \times 1 = 1$   
 $(1 \times \omega) \times \omega^2 = \omega \times \omega^2 = 1$
- iii) Since  $1 \times 1 = 1$ ,  $1 \times \omega = \omega \times 1 = \omega$ ,  $1 \times \omega^2 = \omega^2 \times 1 = \omega^2$   
 Thus 1 is an identity element in  $G$ .
- iv) Since  $1 \times 1 = 1 \times 1 = 1$ ,  $\omega \times \omega^2 = \omega^2 \times \omega = 1$ ,  $\omega^2 \times \omega = \omega \times \omega^2 = 1$   
 therefore inverse of 1 is 1, inverse of  $\omega$  is  $\omega^2$ , inverse of  $\omega^2$  is  $\omega$ .
- v) As table is symmetric about principle diagonal therefore commutative law holds in  $G$ .

Hence  $G$  is an abelian group under multiplication.

### Question # 6

If  $G$  is a group under the operation  $*$  and  $a, b \in G$ , find the solutions of the equations:  $a * x = b$ ,  $x * a = b$

#### Solution

Given that  $G$  is a group under the operation  $*$  and  $a, b \in G$  such that

$$a * x = b$$

As  $a \in G$  and  $G$  is group so  $a^{-1} \in G$  such that

$$a^{-1} * (a * x) = a^{-1} * b$$

$$\Rightarrow (a^{-1} * a) * x = a^{-1} * b \quad \text{as associative law hold in } G.$$

$$\Rightarrow e * x = a^{-1} * b \quad \text{by inverse law.}$$

$$\Rightarrow x = a^{-1} * b \quad \text{by identity law.}$$

And for

$$x * a = b$$

$$\Rightarrow (x * a) * a^{-1} = b * a^{-1} \quad \text{For } a \in G, a^{-1} \in G$$

$$\Rightarrow x * (a * a^{-1}) = b * a^{-1} \quad \text{as associative law hold in } G.$$

$$\Rightarrow x * e = b * a^{-1} \quad \text{by inverse law.}$$

$$\Rightarrow x = b * a^{-1} \quad \text{by identity law.}$$

### Question # 7

Show that the set consisting of elements of the form  $a + \sqrt{3}b$  ( $a, b$  being rational), is an abelian group w.r.t. addition.

#### Solution

$$\text{Consider } G = \{a + \sqrt{3}b \mid a, b \in \mathbb{Q}\}$$

- i) Let  $a + \sqrt{3}b, c + \sqrt{3}d \in G$ , where  $a, b, c$  &  $d$  are rational.

$$(a + \sqrt{3}b) + (c + \sqrt{3}d) = (a + c) + \sqrt{3}(b + d) = a' + \sqrt{3}b' \in G$$

where  $a' = a + c$  and  $b' = b + d$  are rational as sum of rational is rational.

Thus closure law holds in  $G$  under addition.

- ii) For  $a + \sqrt{3}b, c + \sqrt{3}d, e + \sqrt{3}f \in G$

$$\begin{aligned}
(a + \sqrt{3}b) + ((c + \sqrt{3}d) + (e + \sqrt{3}f)) &= (a + \sqrt{3}b) + ((c + e) + \sqrt{3}(d + f)) \\
&= (a + (c + e)) + \sqrt{3}(b + (d + f)) \\
&= ((a + c) + e) + \sqrt{3}((b + d) + f) \\
&\quad \text{As associative law hold in } \mathbb{Q} \\
&= ((a + c) + \sqrt{3}(b + d)) + (e + \sqrt{3}f) \\
&= ((a + \sqrt{3}b) + (c + \sqrt{3}d)) + (e + \sqrt{3}f)
\end{aligned}$$

Thus associative law hold in  $G$  under addition.

iii)  $0 + \sqrt{3} \cdot 0 \in G$  as 0 is a rational such that for any  $a + \sqrt{3}b \in G$

$$(a + \sqrt{3}b) + (0 + \sqrt{3} \cdot 0) = (a + 0) + \sqrt{3}(b + 0) = a + \sqrt{3}b$$

$$\text{And } (0 + \sqrt{3} \cdot 0) + (a + \sqrt{3}b) = (0 + a) + \sqrt{3}(0 + b) = a + \sqrt{3}b$$

Thus  $0 + \sqrt{3} \cdot 0$  is an identity element in  $G$ .

iv) For  $a + \sqrt{3}b \in G$  where  $a$  &  $b$  are rational there exist rational  $-a$  &  $-b$  such that

$$(a + \sqrt{3}b) + ((-a) + \sqrt{3}(-b)) = (a + (-a)) + \sqrt{3}(b + (-b)) = 0 + \sqrt{3} \cdot 0$$

$$\& ((-a) + \sqrt{3}(-b)) + (a + \sqrt{3}b) = ((-a) + a) + \sqrt{3}((-b) + b) = 0 + \sqrt{3} \cdot 0$$

Thus inverse of  $a + \sqrt{3}b$  is  $(-a) + \sqrt{3}(-b)$  exists in  $G$ .

v) For  $a + \sqrt{3}b, c + \sqrt{3}d \in G$

$$\begin{aligned}
(a + \sqrt{3}b) + (c + \sqrt{3}d) &= (a + c) + \sqrt{3}(b + d) \\
&= (c + a) + \sqrt{3}(d + b) \quad \text{As commutative law hold in } \mathbb{Q}. \\
&= (c + d\sqrt{3}) + (a + \sqrt{3}b)
\end{aligned}$$

Thus Commutative law holds in  $G$  under addition.

And hence  $G$  is an abelian group under addition.

### Question 8

Determine whether  $(P(S), *)$ , where  $*$  stands for intersection is a semi group, a monoid or neither. If it is a monoid, specify its identity.

#### Solution

Let  $A, B \in P(S)$  where  $A$  &  $B$  are subsets of  $S$ .

As intersection of two subsets of  $S$  is subset of  $S$ .

Therefore  $A * B = A \cap B \in P(S)$ . Thus closure law holds in  $P(S)$ .

For  $A, B, C \in P(S)$

$$A * (B * C) = A \cap (B \cap C) = (A \cap B) \cap C = (A * B) * C$$

Thus associative law holds and  $P(S)$ .

And hence  $(P(S), *)$  is a semi-group.

For  $A \in P(S)$  where  $A$  is a subset of  $S$  we have  $S \in P(S)$  such that

$$A \cap S = S \cap A = A.$$

Thus  $S$  is an identity element in  $P(S)$ . And hence  $(P(S), *)$  is a monoid.

**Question 9**

Complete the following table to obtain a semi-group under \*

**Solution**

Let  $x_1$  and  $x_2$  be the required elements.

By associative law

$$(a * a) * a = a * (a * a)$$

$$\Rightarrow c * a = a * c$$

$$\Rightarrow x_1 = b$$

Now again by associative law

$$(a * a) * b = a * (a * b)$$

$$\Rightarrow c * b = a * a \Rightarrow x_2 = c$$

*	a	b	c
a	c	a	b
b	a	b	c
c	$x_1$	$x_2$	a

**Question 10**

Prove that all  $2 \times 2$  non-singular matrices over the real field form a non-abelian group under multiplication.

**Solution** Let  $G$  be the all non-singular  $2 \times 2$  matrices over the real field.

i) Let  $A, B \in G$  then  $A_{2 \times 2} \times B_{2 \times 2} = C_{2 \times 2} \in G$

Thus closure law holds in  $G$  under multiplication.

ii) Associative law in matrices of same order under multiplication holds.

therefore for  $A, B, C \in G$

$$A \times (B \times C) = (A \times B) \times C$$

iii)  $I_{2 \times 2} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$  is a non-singular matrix such that

$$A_{2 \times 2} \times I_{2 \times 2} = I_{2 \times 2} \times A_{2 \times 2} = A_{2 \times 2}$$

Thus  $I_{2 \times 2}$  is an identity element in  $G$ .

iv) Since inverse of non-singular square matrix exists,

therefore for  $A \in G$  there exist  $A^{-1} \in G$  such that  $AA^{-1} = A^{-1}A = I$ .

v) As we know for any two matrices  $A, B \in G$ ,  $AB \neq BA$  in general.

Therefore commutative law does not hold in  $G$  under multiplication.

Hence the set of all  $2 \times 2$  non-singular matrices over a real field is a non-abelian group under multiplication.