

Q No. 1 i) $a_1 = 5$ and other three consecutive terms are 23, 26, 29.
 Since $a_1 = 5$ & $d = 26 - 23 = 3$.
 Now, $a_2 = a_1 + d = 5 + 3 = 8$
 $a_3 = a_2 + d = 8 + 3 = 11$
 $a_4 = a_3 + d = 11 + 3 = 14$.
 hence 5, 8, 11, 14 are first four terms of A.P.

Q No. 2 (i) $a_5 = 17$ and $a_9 = 37$
 Consider a_1 be the first term and 'd' be the common difference.

$$\text{Since } a_5 = 17 \\ \Rightarrow a_1 + (5-1)d = 17 \\ \Rightarrow a_1 + 4d = 17 \quad \text{(i)}$$

$$\text{also } a_9 = 37 \\ \Rightarrow a_1 + (9-1)d = 37 \\ \Rightarrow a_1 + 8d = 37 \quad \text{(ii)}$$

Subtracting (i) and (ii)

$$\begin{array}{r} a_1 + 4d = 17 \\ a_1 + 8d = 37 \\ \hline -4d = -20 \\ \Rightarrow d = 5 \end{array}$$

putting value of d in (i)

$$a_1 + 4(5) = 17$$

$$\Rightarrow a_1 + 20 = 17$$

$$\Rightarrow a_1 = 17 - 20 \quad \cancel{\text{cancel}}$$

$$\Rightarrow a_1 = -3$$

$$\text{So } a_2 = a_1 + d = -3 + 5 = 2$$

$$a_3 = a_2 + d = 2 + 5 = 7$$

$$a_4 = a_3 + d = 7 + 5 = 12$$

hence -3, 2, 7, 12 are first four terms of A.P.

$$\text{iii) } 3a_7 = 7a_4 \& a_{10} = 33$$

Suppose a_1 be the first term and d be the common difference.

Since $3a_7 = 7a_4$

$$\Rightarrow 3(a_1 + 6d) = 7(a_1 + 3d)$$

$$\Rightarrow 3a_1 + 18d = 7a_1 + 21d$$

$$\Rightarrow 3a_1 + 18d - 7a_1 - 21d = 0$$

$$\Rightarrow -4a_1 - 3d = 0$$

$$\Rightarrow 4a_1 + 3d = 0 \quad \text{(i)}$$

also $a_{10} = 33$

$$\Rightarrow a_1 + 9d = 33 \quad \text{(ii)}$$

cross eq. (i) by 4 & subtracting from (ii)

$$\begin{array}{r} 4a_1 + 3d = 0 \\ 4a_1 + 36d = 132 \\ \hline -33d = -132 \\ \Rightarrow d = \frac{-132}{-33} = 4 \end{array}$$

putting value of d in (ii)

$$a_1 + 9(4) = 33$$

$$\Rightarrow a_1 + 36 = 33$$

$$\Rightarrow a_1 = 33 - 36 \Rightarrow a_1 = -3$$

$$\text{So } a_2 = a_1 + d = -3 + 4 = 1$$

$$a_3 = a_2 + d = 1 + 4 = 5$$

$$a_4 = a_3 + d = 5 + 4 = 9$$

hence -3, 1, 5, 9 are the first four terms of A.P.

Q No. 2 $a_{n-3} = 2n - 5$

$$\Rightarrow a_{n-3} = 2n - 6 + 1$$

$$= 2(n-3) + 1$$

Replacing $n-3$ by n .

$$a_n = 2n + 1$$

Answer

QNo3 Suppose a_1 be the first term and d be common difference of A.P.

$$\text{Since } a_5 = 16$$

$$\Rightarrow a_1 + 4d = 16 \quad \text{(i)}$$

$$\text{also } a_{20} = 46$$

$$\Rightarrow a_1 + 19d = 46 \quad \text{(ii)}$$

Subtracting (i) & (ii)

$$\cancel{a_1 + 4d = 16}$$

$$\cancel{a_1 + 19d = 46}$$

$$= 15d = -30$$

$$\Rightarrow d = -2$$

putting value of d in (i)

$$a_1 + 4(2) = 16$$

$$\Rightarrow a_1 + 8 = 16$$

$$\Rightarrow a_1 = 16 - 8 \Rightarrow a_1 = 8$$

Now

$$a_{12} = a_1 + 11d$$

$$= 8 + 11(-2)$$

$$= 8 + 22 = 30$$

Answer

QNo.4

$$x, 1, 2-x, 3-2x, \dots$$

$$\text{here } a_1 = 1$$

$$\text{and } d = a_2 - a_1$$

$$= 1 - x$$

Since

$$a_{13} = a_1 + 12d$$

$$= x + 12(1-x)$$

$$= x + 12 - 12x$$

~~$\therefore a_{13} = a_1 + 12d$~~

$$\therefore a_{13} = 12 - 11x \quad \text{Answer}$$

QNo5 Same as QNo.3

QNo.6

$$5, 2, -1, \dots, -85$$

$$\text{here } a_1 = 5$$

$$d = a_2 - a_1 = 2 - 5 = -3$$

$$a_n = -85, n = ?$$

Since

$$a_n = a_1 + (n-1)d$$

$$\Rightarrow -85 = 5 + (n-1)(-3)$$

$$\Rightarrow -85 = 5 - 3n + 3$$

$$\Rightarrow 3n = 5 + 3 + 85$$

$$\Rightarrow 3n = 93$$

$$\Rightarrow \boxed{n = 31} \quad \text{Answer}$$

QNo.7 Same as above

QNo.8 $a_1 = 11, a_n = 68$

$$d = 3, n = ?$$

Since

$$a_n = a_1 + (n-1)d$$

$$\Rightarrow 68 = 11 + (n-1)3$$

Now solve yourself as above

QNo.9

$$\text{Since } a_n = 3n - 1$$

$$\text{put } n = 1$$

$$a_1 = 3(1) - 1 = 3 - 1 = 2$$

$$\text{put } n = 2$$

$$a_2 = 3(2) - 1 = 6 - 1 = 5$$

$$\text{put } n = 3$$

$$a_3 = 3(3) - 1 = 9 - 1 = 8$$

$$\text{put } n = 4$$

$$a_4 = 3(4) - 1 = 12 - 1 = 11$$

Thus

$$2, 5, 8, 11, \dots$$

is the required A.P

(Q) No 10. 17, 13, 9, ... , ... , ...

$$a_1 = 17, d = 13 - 17 = -4$$

i) Suppose -19 be the n th term of A.P. i.e. $a_n = -19$

$$\text{Since } a_n = a_1 + (n-1)d$$

$$\Rightarrow -19 = 17 + (n-1)(-4)$$

$$\Rightarrow -19 = 17 - 4n + 4$$

$$\Rightarrow 4n = 17 + 4 + 19$$

$$= 40$$

$$\Rightarrow n = 10$$

Thus -19 is the 10th term of A.P.

ii) Suppose 2 be the n th term of A.P. i.e. $a_n = 2$

$$\text{Since } a_n = a_1 + (n-1)d$$

$$\Rightarrow 2 = 17 + (n-1)(-4)$$

$$\Rightarrow 2 = 17 - 4n + 4$$

$$\Rightarrow 4n = 17 + 4 - 2$$

$$= 19$$

$\Rightarrow n = \frac{19}{4}$ which is a rational, therefore 2 is not the term of A.P.

(Q) No 11.

Let a_1 be the first term and d be the common difference

$$\text{Now } a_p = l$$

$$\Rightarrow a_1 + (p-1)d = l$$

$$\therefore a_q = m$$

$$\Rightarrow a_1 + (q-1)d = m$$

$$\therefore a_r = n$$

$$\Rightarrow a_1 + (r-1)d = n$$

$$i) L.H.S = l(q-r) + m(r-p) + n(p-q)$$

$$= [a_1 + (p-1)d](q-r) + [a_1 + (q-1)d](r-p)$$

$$+ [a_1 + (r-1)d](p-q)$$

$$= (a_1 + pd - d)(q-r) + (a_1 + qd - q)(r-p)$$

$$+ (a_1 + rd - r)(p-q)$$

$$= a_1q + pdqd - qd - 2a_1r + pd + rd + pq$$

$$+ a_1p + prd - pd - 3a_1q - qrd + qd$$

$$= 0 = R.H.S \quad \text{proved}$$

$$ii) L.H.S = p(m-n) + q(n-l) + r(l-m)$$

$$= p[a_1 + (q-1)d - a_1 - (r-1)d]$$

$$+ q[a_1 + (r-1)d - a_1 - (p-1)d]$$

$$= p[qd - d - rd + d]$$

$$+ q[rd - d - pd + d]$$

$$+ r[pd - d - qd + d]$$

$$= pd - pd + qrd - pd$$

$$+ pd - qrd = 0 = R.H.S$$

proved

(Q) No 12.

$$(\frac{4}{3}), (\frac{7}{3}), (\frac{10}{3}), \dots$$

We first find the n th term of

$$4, 7, 10, \dots$$

$$a_1 = 4, d = 7 - 4 = 3$$

$$\text{So } a_n = a_1 + (n-1)d$$

$$= 4 + (n-1)3$$

$$= 4 + 3n - 3 = 3n + 1$$

Hence n th term of

$$\text{given sequence is } (\frac{3n+1}{3})^2$$

Q No 13 Since $\frac{1}{a}, \frac{1}{b}, \frac{1}{c}$ are in A.P. therefore

$$d = \frac{1}{b} - \frac{1}{a} = \frac{1}{c} - \frac{1}{b}$$

$$\Rightarrow \frac{1}{b} + \frac{1}{b} = \frac{1}{a} + \frac{1}{c}$$

$$\Rightarrow \frac{1+1}{b} = \frac{c+a}{ac}$$

$$\Rightarrow \frac{2}{b} = \frac{a+c}{ac}$$

$$\Rightarrow \frac{b}{2} = \frac{ac}{a+c}$$

$$\Rightarrow b = \frac{2ac}{a+c} \text{ proved}$$

Q No 14 Since $\frac{1}{a}, \frac{1}{b}, \frac{1}{c}$ are in A.P. therefore

$$d = \frac{1}{b} - \frac{1}{a} \quad \text{(i)}$$

also

$$d = \frac{1}{c} - \frac{1}{b} \quad \text{(ii)}$$

Comparing (i) and (ii)

$$\frac{1}{b} - \frac{1}{a} = \frac{1}{c} - \frac{1}{b}$$

$$\Rightarrow \frac{1}{b} + \frac{1}{b} = \frac{1}{c} + \frac{1}{a}$$

$$\Rightarrow \frac{1+1}{b} = \frac{a+c}{ac}$$

$$\Rightarrow \frac{2}{b} = \frac{a+c}{ac}$$

$$\Rightarrow \frac{b}{2} = \frac{ac}{a+c}$$

$$\Rightarrow b = \frac{2ac}{a+c}$$

putting value of b in eq. (i)

$$d = \frac{1}{\frac{2ac}{a+c}} - \frac{1}{a} = \frac{a+c}{2ac} - \frac{1}{a}$$

$$= \frac{a+c-2c}{2ac} = \frac{a-c}{2ac}$$

Hence the common difference

is $\frac{a-c}{2ac}$

END.