

# Chapter 8

# Exercise 8.3

## Binomial Theorem when n is negative or fraction:

When n is negative or fraction and  $|x| < 1$  then

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \frac{n(n-1)(n-2)}{3!}x^3 + \dots$$

Where the general term of binomial expansion is

$$T_{r+1} = \frac{n(n-1)(n-2)\dots(n-(r-1))}{r!}x^r$$

## Question # 1

Expand the following upto 4 times, taking the values of  $x$  such that the expansion in each case is valid.

- |   |                                |                                    |                                    |
|---|--------------------------------|------------------------------------|------------------------------------|
| (i) $(1-x)^{\frac{1}{2}}$                 | (ii) $(1+2x)^{-1}$             | (iii) $(1+x)^{\frac{1}{3}}$        | (iv) $(4-3x)^{\frac{1}{2}}$        |
| (v) $(8-2x)^{-1}$                         | (vi) $(2-3x)^{-2}$             | (vii) $\frac{(1-x)^{-1}}{(1+x)^2}$ | (viii) $\frac{\sqrt{1+2x}}{(1-x)}$ |
| (ix) $\frac{(4+2x)^{\frac{1}{2}}}{(2-x)}$ | (x) $(1+x-2x^2)^{\frac{1}{2}}$ | (xi) $(1-2x+3x^2)^{\frac{1}{2}}$   |                                    |

### Solution

$$\begin{aligned} \text{(i)} \quad (1-x)^{\frac{1}{2}} &= 1 + \frac{1}{2}(-x) + \frac{\frac{1}{2}\left(\frac{1}{2}-1\right)}{2!}(-x)^2 + \frac{\frac{1}{2}\left(\frac{1}{2}-1\right)\left(\frac{1}{2}-2\right)}{3!}(-x)^3 + \dots \\ &= 1 - \frac{1}{2}x + \frac{\frac{1}{2}\left(-\frac{1}{2}\right)}{2}x^2 + \frac{\frac{1}{2}\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)}{3 \cdot 2}(-x^3) + \dots \\ &= 1 - \frac{1}{2}x - \frac{1}{8}x^2 - \frac{1}{16}x^3 + \dots \end{aligned}$$

$$\text{(ii)} \quad \text{Do yourself as above}$$

$$\text{(iii)} \quad \text{Do yourself as above}$$

$$\text{(iv)} \quad (4-3x)^{\frac{1}{2}} = \left[ 4\left(1-\frac{3x}{4}\right) \right]^{\frac{1}{2}} = (4)^{\frac{1}{2}} \left(1-\frac{3x}{4}\right)^{\frac{1}{2}} = 2\left(1-\frac{3x}{4}\right)^{\frac{1}{2}}$$

$$= 2 \left[ 1 + \frac{1}{2}\left(-\frac{3x}{4}\right) + \frac{\frac{1}{2}\left(\frac{1}{2}-1\right)}{2!} \left(-\frac{3x}{4}\right)^2 + \frac{\frac{1}{2}\left(\frac{1}{2}-1\right)\left(\frac{1}{2}-2\right)}{3!} \left(-\frac{3x}{4}\right)^3 + \dots \right]$$

$$\begin{aligned}
 &= 2 \left[ 1 - \frac{3x}{8} + \frac{1}{2} \left( -\frac{1}{2} \right) \left( \frac{9x^2}{16} \right) + \frac{1}{2} \left( -\frac{1}{2} \right) \left( -\frac{3}{2} \right) \left( -\frac{27x^3}{64} \right) + \dots \right] \\
 &= 2 \left[ 1 - \frac{3x}{8} - \frac{1}{8} \left( \frac{9x^2}{16} \right) - \frac{1}{16} \left( \frac{27x^3}{64} \right) + \dots \right] \\
 &= 2 \left[ 1 - \frac{3x}{8} - \frac{9x^2}{128} - \frac{27x^3}{1024} + \dots \right] \\
 &= 2 - \frac{3x}{4} - \frac{9x^2}{64} - \frac{27x^3}{512} + \dots
 \end{aligned}$$

(v)  $(8 - 2x)^{\frac{1}{2}} = (8)^{-1} \left( 1 - \frac{2x}{8} \right)^{-1} = \frac{1}{8} \left( 1 - \frac{x}{4} \right)^{-1}$  Now do yourself

(vi) Do yourself

(vii)  $\frac{(1-x)^{-1}}{(1+x)^2} = (1-x)^{-1}(1+x)^{-2}$

$$\begin{aligned}
 &= \left( 1 + (-1)(-x) + \frac{(-1)(-1-1)}{2!}(-x)^2 + \frac{(-1)(-1-1)(-1-2)}{3!}(-x)^3 + \dots \right) \\
 &\quad \times \left( 1 + (-2)(x) + \frac{(-2)(-2-1)}{2!}(x)^2 + \frac{(-2)(-2-1)(-2-2)}{3!}(x)^3 + \dots \right) \\
 &= \left( 1 + x + \frac{(-1)(-2)}{2}(x^2) + \frac{(-1)(-2)(-3)}{3 \cdot 2}(-x^3) + \dots \right) \\
 &\quad \times \left( 1 - 2x + \frac{(-2)(-3)}{2}(x)^2 + \frac{(-2)(-3)(-4)}{3 \cdot 2}(x)^3 + \dots \right) \\
 &= (1 + x + x^2 + x^3 + \dots) \times (1 - 2x + 3x^2 - 4x^3 + \dots) \\
 &= 1 + (x - 2x) + (x^2 - 2x^2 + 3x^2) + (x^3 - 2x^3 + 3x^3 - 4x^3) + \dots \\
 &= 1 - x + 2x^2 - 2x^3 + \dots
 \end{aligned}$$

(viii) Do yourself as above

(ix)  $\frac{(4+2x)^{\frac{1}{2}}}{2-x} = (4+2x)^{\frac{1}{2}}(2-x)^{-1} = (4)^{\frac{1}{2}} \left( 1 + \frac{2x}{4} \right)^{\frac{1}{2}} (2)^{-1} \left( 1 - \frac{x}{2} \right)^{-1}$

$$\begin{aligned}
 &= (4)^{\frac{1}{2}} \left( 1 + \frac{x}{2} \right)^{\frac{1}{2}} (2)^{-1} \left( 1 - \frac{x}{2} \right)^{-1} = 2 \left( 1 + \frac{x}{2} \right)^{\frac{1}{2}} \frac{1}{2} \left( 1 - \frac{x}{2} \right)^{-1} = \left( 1 + \frac{x}{2} \right)^{\frac{1}{2}} \left( 1 - \frac{x}{2} \right)^{-1}
 \end{aligned}$$

$$\begin{aligned}
&= \left(1 + \frac{x}{2}\right)^{\frac{1}{2}} \left(1 - \frac{x}{2}\right)^{-1} \\
&= \left(1 + \frac{1}{2} \left(\frac{x}{2}\right) + \frac{\frac{1}{2} \left(\frac{1}{2}-1\right)}{2!} \left(\frac{x}{2}\right)^2 + \frac{\frac{1}{2} \left(\frac{1}{2}-1\right) \left(\frac{1}{2}-2\right)}{3!} \left(\frac{x}{2}\right)^3 + \dots\right) \\
&\quad \times \left(1 + (-1) \left(-\frac{x}{2}\right) + \frac{(-1)(-1-1)}{2!} \left(-\frac{x}{2}\right)^2 + \frac{(-1)(-1-1)(-1-2)}{3!} \left(-\frac{x}{2}\right)^3 + \dots\right) \\
&= \left(1 + \frac{x}{4} + \frac{\frac{1}{2} \left(-\frac{1}{2}\right)}{2} \left(\frac{x^2}{4}\right) + \frac{\frac{1}{2} \left(-\frac{1}{2}\right) \left(-\frac{3}{2}\right)}{3 \cdot 2} \left(\frac{x^3}{8}\right) + \dots\right) \\
&\quad \times \left(1 + \frac{x}{2} + \frac{(-1)(-2)}{2} \left(\frac{x^2}{4}\right) + \frac{(-1)(-2)(-3)}{3 \cdot 2} \left(-\frac{x^3}{8}\right) + \dots\right) \\
&= \left(1 + \frac{x}{4} - \frac{x^2}{32} + \frac{x^3}{128} + \dots\right) \times \left(1 + \frac{x}{2} + \frac{x^2}{4} + \frac{x^3}{8} + \dots\right) \\
&= 1 + \left(\frac{x}{4} + \frac{x}{2}\right) + \left(-\frac{x^2}{32} + \frac{x^2}{8} + \frac{x^2}{4}\right) + \left(\frac{x^3}{128} - \frac{x^3}{64} + \frac{x^3}{16} + \frac{x^3}{8}\right) + \dots \\
&= 1 + \frac{3x}{4} + \frac{11x^2}{32} + \frac{23x^3}{128} + \dots
\end{aligned}$$

$$\begin{aligned}
(x) \quad & (1+x-2x^2)^{\frac{1}{2}} = (1+(x-2x^2))^{\frac{1}{2}} \\
&= 1 + \frac{1}{2}(x-2x^2) + \frac{\frac{1}{2} \left(\frac{1}{2}-1\right)}{2!} (x-2x^2)^2 + \frac{\frac{1}{2} \left(\frac{1}{2}-1\right) \left(\frac{1}{2}-2\right)}{3!} (x-2x^2)^3 + \dots \\
&= 1 + \frac{1}{2}(x-2x^2) + \frac{\frac{1}{2} \left(-\frac{1}{2}\right)}{2} (x^2 - 4x^3 + 4x^4) + \frac{\frac{1}{2} \left(-\frac{1}{2}\right) \left(-\frac{3}{2}\right)}{3 \cdot 2} \\
&\quad (x^3 + 3(x)^2(-2x^2) + 3(x)(-2x^2)^2 - (2x^2)^3) + \dots \\
&= 1 + \frac{1}{2}(x-2x^2) - \frac{1}{8}(x^2 - 4x^3 + 4x^4) + \frac{1}{16}(x^3 - 6x^4 + 12x^5 - 8x^6) + \dots \\
&= 1 + \frac{1}{2}x - \frac{2}{2}x^2 - \frac{1}{8}x^2 - \frac{4}{8}x^3 + \frac{4}{8}x^4 + \frac{1}{16}x^3 - \frac{6}{16}x^4 + \frac{12}{16}x^5 - \frac{8}{16}x^6 + \dots \\
&= 1 + \frac{1}{2}x - x^2 - \frac{1}{8}x^2 - \frac{1}{2}x^3 + \frac{1}{2}x^4 + \frac{1}{16}x^3 - \frac{3}{8}x^4 + \frac{3}{4}x^5 - \frac{1}{8}x^6 + \dots \\
&= 1 + \frac{1}{2}x - \frac{9}{8}x^2 - \frac{9}{16}x^3 + \dots
\end{aligned}$$

(xi)

*Do yourself as above*

**Question # 2**

Use the Binomial theorem find the value of the following to three places of decimals.

(i)  $\sqrt{99}$

(ii)  $(0.98)^{\frac{1}{2}}$

(iii)  $(1.03)^{\frac{1}{3}}$

(iv)  $\sqrt[3]{65}$

(v)  $\sqrt[4]{17}$

(vi)  $\sqrt[4]{31}$

(vii)  $\frac{1}{\sqrt[3]{998}}$

(viii)  $\frac{1}{\sqrt[5]{252}}$

(ix)  $\frac{\sqrt{7}}{\sqrt{8}}$

(x)  $(0.998)^{\frac{1}{3}}$

(xi)  $\frac{1}{\sqrt[6]{486}}$

(xii)  $(1280)^{\frac{1}{4}}$

**Solution**

$$\begin{aligned}
 \text{(i)} \quad \sqrt{99} &= (99)^{\frac{1}{2}} = (100-1)^{\frac{1}{2}} = (100)^{\frac{1}{2}} \left(1 - \frac{1}{100}\right)^{\frac{1}{2}} \\
 &= 10 \left(1 + \frac{1}{2} \left(-\frac{1}{100}\right) + \frac{\frac{1}{2} \left(\frac{1}{2}-1\right)}{2!} \left(-\frac{1}{100}\right)^2 + \dots\right) \\
 &= 10 \left(1 - \frac{1}{200} + \frac{\frac{1}{2} \left(-\frac{1}{2}\right)}{2} \left(\frac{1}{10000}\right) + \dots\right) \\
 &= 10 \left(1 - 0.005 - \frac{1}{8}(0.0001) + \dots\right) \\
 &= 10(1 - 0.005 - 0.0000125 + \dots) \\
 &\approx 10(0.9949875) = 9.949875 \\
 &\approx 9.950
 \end{aligned}$$

$$\text{(ii)} \quad (0.98)^{\frac{1}{2}} = (1-0.02)^{\frac{1}{2}} \quad \text{Now do yourself}$$

$$\text{(iii)} \quad (1.03)^{\frac{1}{3}} = (1+0.03)^{\frac{1}{3}} \quad \text{Now do yourself}$$

$$\text{(iv)} \quad \sqrt[3]{65} = (65)^{\frac{1}{3}} = (64-1)^{\frac{1}{3}} = (64)^{\frac{1}{3}} \left(1 - \frac{1}{64}\right)^{\frac{1}{3}} \quad \text{Now do yourself}$$

$$\text{(v)} \quad \sqrt[4]{17} = (17)^{\frac{1}{4}} = (16-1)^{\frac{1}{4}} = (16)^{\frac{1}{4}} \left(1 - \frac{1}{16}\right)^{\frac{1}{4}} \quad \text{Now do yourself}$$

$$\text{(vi)} \quad \sqrt[5]{31} = (31)^{\frac{1}{5}} = (32-1)^{\frac{1}{5}} = (32)^{\frac{1}{5}} \left(1 - \frac{1}{32}\right)^{\frac{1}{5}} \quad \text{Now do yourself}$$

$$\begin{aligned}
 \text{(vii)} \quad \frac{1}{\sqrt[3]{998}} &= \frac{1}{(998)^{\frac{1}{3}}} = (998)^{-\frac{1}{3}} = (1000 - 2)^{-\frac{1}{3}} = (1000)^{-\frac{1}{3}} \left(1 - \frac{2}{1000}\right)^{-\frac{1}{3}} \\
 &= (10^3)^{-\frac{1}{3}} \left(1 - \frac{1}{500}\right)^{-\frac{1}{3}} \\
 &= \left(\frac{1}{10}\right) \left(1 + \left(-\frac{1}{3}\right) \left(-\frac{1}{500}\right) + \frac{-\frac{1}{3} \left(-\frac{1}{3} - 1\right)}{2!} \left(-\frac{1}{500}\right)^2 + \dots\right) \\
 &= \left(\frac{1}{10}\right) \left(1 + \left(\frac{1}{1500}\right) + \frac{-\frac{1}{3} \left(-\frac{4}{3}\right)}{2} \left(\frac{1}{250000}\right) + \dots\right) \\
 &= \left(\frac{1}{10}\right) \left(1 + (0.0006667) + \frac{2}{9} (0.000004) + \dots\right) \\
 &= \left(\frac{1}{10}\right) (1 + 0.0006667 + 0.00000089 + \dots) \\
 &\approx \left(\frac{1}{10}\right) (1.00066759) = 0.100066759 \approx 0.100 \quad \text{Answer}
 \end{aligned}$$

$$\begin{aligned}
 \text{(viii)} \quad \frac{1}{\sqrt[5]{252}} &= \frac{1}{(252)^{\frac{1}{5}}} = (252)^{-\frac{1}{5}} = (243 + 9)^{-\frac{1}{5}} = (243)^{\frac{1}{5}} \left(1 + \frac{9}{243}\right)^{-\frac{1}{5}} \\
 &= (3^5)^{-\frac{1}{5}} \left(1 + \frac{1}{27}\right)^{-\frac{1}{5}} \quad \text{Now do yourself as above}
 \end{aligned}$$

$$\begin{aligned}
 \text{(ix)} \quad \frac{\sqrt{7}}{\sqrt{8}} &= \sqrt{\frac{7}{8}} = \left(\frac{7}{8}\right)^{\frac{1}{2}} = \left(1 - \frac{1}{8}\right)^{\frac{1}{2}} \\
 &= 1 + \frac{1}{2} \left(-\frac{1}{8}\right) + \frac{\frac{1}{2} \left(\frac{1}{2} - 1\right)}{2!} \left(-\frac{1}{8}\right)^2 + \dots \\
 &= 1 - \frac{1}{16} + \frac{\frac{1}{2} \left(-\frac{1}{2}\right)}{2} \left(\frac{1}{64}\right) + \dots \\
 &= 1 - \frac{1}{16} - \frac{1}{8} \left(\frac{1}{64}\right) + \dots \\
 &= 1 - \frac{1}{16} - \frac{1}{512} + \dots \\
 &= 1 - 0.0625 - 0.00195 + \dots \\
 &\approx 0.93555 \approx 0.936 \quad \text{Answer}
 \end{aligned}$$

$$(x) \quad (0.998)^{-\frac{1}{3}} = (1 - 0.002)^{-\frac{1}{3}} \quad \text{Now do yourself as above}$$

$$\begin{aligned} (xi) \quad \frac{1}{\sqrt[6]{486}} &= \frac{1}{(486)^{\frac{1}{6}}} = (486)^{-\frac{1}{6}} = (729 - 243)^{-\frac{1}{6}} = (729)^{\frac{1}{6}} \left(1 - \frac{243}{729}\right)^{-\frac{1}{6}} \\ &= (3^6)^{-\frac{1}{6}} \left(1 - \frac{1}{3}\right)^{-\frac{1}{6}} \quad \text{Now do yourself} \end{aligned}$$

$$\begin{aligned} (xii) \quad (1280)^{\frac{1}{4}} &= (1296 - 16)^{\frac{1}{4}} = (1296)^{\frac{1}{4}} \left(1 - \frac{16}{1296}\right)^{\frac{1}{4}} = (6^4)^{\frac{1}{4}} \left(1 - \frac{1}{81}\right)^{\frac{1}{4}} \\ &\quad \text{Now do yourself} \end{aligned}$$


---

### Question # 3

Find the coefficient of  $x^n$  in the expansion of

- |                                |                                |                                 |
|--------------------------------|--------------------------------|---------------------------------|
| (i) $\frac{1+x^2}{(1+x)^2}$    | (ii) $\frac{(1+x)^2}{(1-x)^2}$ | (iii) $\frac{(1+x)^3}{(1-x)^2}$ |
| (iv) $\frac{(1+x)^2}{(1-x)^3}$ | (v) $(1-x+x^2-x^3+\dots)^2$    |                                 |

#### Solution

$$\begin{aligned} (i) \quad \frac{1+x^2}{(1+x)^2} &= (1+x^2)(1+x)^{-2} \\ &= (1+x^2) \left(1 + (-2)(x) + \frac{(-2)(-2-1)}{2!}(x)^2 + \frac{(-2)(-2-1)(-2-2)}{3!}(x)^3 + \dots\right) \\ &= (1+x^2) \left(1 - 2x + \frac{(-2)(-3)}{2}x^2 + \frac{(-2)(-3)(-4)}{3 \cdot 2}x^3 + \dots\right) \\ &= (1+x^2)(1 - 2x + 3x^2 - 4x^3 + \dots) \\ &= (1+x^2)(1 + (-1)2x + (-1)^23x^2 + (-1)^34x^3 + \dots) \end{aligned}$$

Following in this way we can write

$$\begin{aligned} \frac{1+x^2}{(1+x)^2} &= (1+x^2)(1 + (-1)2x + (-1)^23x^2 + (-1)^34x^3 + \dots + (-1)^{n-2}(n-1)x^{n-2} + \\ &\quad (-1)^{n-1}(n)x^{n-1} + (-1)^n(n+1)x^n + \dots) \end{aligned}$$

So taking only terms involving  $x^n$  we get

$$\begin{aligned} &(-1)^n(n+1)x^n + (-1)^{n-2}(n-1)x^n \\ &= (-1)^n(n+1)x^n + (-1)^n(-1)^{-2}(n-1)x^n \\ &= (-1)^n(n+1)x^n + (-1)^n(n-1)x^n \quad \because (-1)^{-2} = 1 \\ &= (n+1+n-1)(-1)^n x^n = (2n)(-1)^n x^n \end{aligned}$$

Thus the coefficient of term involving  $x^n$  is  $(2n)(-1)^n$

(ii)

Hint:

After solving you will get

$$\frac{(1+x)^2}{(1-x)^2} = (1+2x+x^2)(1+2x+3x^2+4x^3+\dots+(n-1)x^{n-2}+(n)x^{n-1}+(n+1)x^n+\dots)$$

*Do yourself as above*

$$\begin{aligned} \text{(iii)} \quad & \frac{(1+x)^3}{(1-x)^2} = (1+x)^3(1-x)^{-2} \\ &= (1+x)^3 \left( 1 + (-2)(-x) + \frac{(-2)(-2-1)}{2!}(-x)^2 + \frac{(-2)(-2-1)(-2-2)}{3!}(-x)^3 + \dots \right) \\ &= (1+x)^3 \left( 1 + 2x + \frac{(-2)(-3)}{2}(-x)^2 + \frac{(-2)(-3)(-4)}{3 \cdot 2}(-x)^3 + \dots \right) \\ &= (1+3x+3x^2+x^3)(1+2x+3x^2+4x^3+\dots) \end{aligned}$$

Following in this way we can write

$$\begin{aligned} \frac{(1+x)^3}{(1-x)^2} &= (1+3x+3x^2+x^3)(1+2x+3x^2+4x^3+\dots+(n-2)x^{n-3}+(n-1)x^{n-2} \\ &\quad +(n)x^{n-1}+(n+1)x^n+\dots) \end{aligned}$$

So taking only terms involving  $x^n$  we have term

$$\begin{aligned} & (n+1)x^n + 3(n)x^n + 3(n-1)x^n + (n-2)x^n \\ &= ((n+1) + 3(n) + 3(n-1) + (n-2))x^n \\ &= (n+1 + 3n + 3n-3 + n-2)x^n \\ &= (8n-4)x^n \end{aligned}$$

Thus the coefficient of term involving  $x^n$  is  $(8n-4)$ .

$$\begin{aligned} \text{(iv)} \quad & \frac{(1+x)^2}{(1-x)^3} = (1+x)^2(1-x)^{-3} \\ &= (1+x)^2 \left( 1 + (-3)(-x) + \frac{(-3)(-3-1)}{2!}(-x)^2 + \frac{(-3)(-3-1)(-3-2)}{3!}(-x)^3 + \dots \right) \\ &= (1+x)^2 \left( 1 + (-3)(-x) + \frac{(-3)(-4)}{2}(-x)^2 + \frac{(-3)(-4)(-5)}{3 \cdot 2}(-x)^3 + \dots \right) \\ &= (1+2x+x^2) \left( 1 + 3x + \frac{(3)(4)}{2}(x^2) + \frac{(4)(5)}{2}(x^3) + \dots \right) \\ &= (1+2x+x^2) \left( 1 + \frac{(2)(3)}{2}x + \frac{(3)(4)}{2}x^2 + \frac{(4)(5)}{2}x^3 + \dots \right) \end{aligned}$$

Following in this way we can write

$$\frac{(1+x)^2}{(1-x)^3} = (1+2x+x^2) \left( 1 + \frac{(2)(3)}{2}x + \frac{(3)(4)}{2}x^2 + \frac{(4)(5)}{2}x^3 + \dots \right)$$

$$+ \frac{(n-1)(n)}{2}x^{n-2} + \frac{(n)(n+1)}{2}x^{n-1} + \frac{(n+1)(n+2)}{2}x^n + \dots \Big)$$

So taking only terms involving  $x^n$  we have term

$$\begin{aligned} & \frac{(n+1)(n+2)}{2}x^n + 2\frac{(n)(n+1)}{2}x^n + \frac{(n-1)(n)}{2}x^n \\ &= ((n+1)(n+2) + 2(n)(n+1) + (n-1)(n)) \frac{x^n}{2} \\ &= (n^2 + n + 2n + 2 + 2n^2 + 2n + n^2 - n) \frac{x^n}{2} \\ &= (4n^2 + 4n + 2) \frac{x^n}{2} = 2(2n^2 + 2n + 1) \frac{x^n}{2} \\ &= (2n^2 + 2n + 1)x^n \end{aligned}$$

Thus the coefficient of term involving  $x^n$  is  $(2n^2 + 2n + 1)$ .

(v) Since we know that

$$(1+x)^{-1} = 1 - x + x^2 - x^3 + \dots$$

Therefore

$$\begin{aligned} (1-x+x^2-x^3+\dots)^2 &= ((1+x)^{-1})^2 = (1+x)^{-2} \\ &= 1 + (-2)(x) + \frac{(-2)(-2-1)}{2!}(x)^2 + \frac{(-2)(-2-1)(-2-2)}{3!}(x)^3 + \dots \\ &= 1 - 2x + \frac{(-2)(-3)}{2}(x)^2 + \frac{(-2)(-3)(-4)}{3 \cdot 2}(x)^3 + \dots \\ &= 1 - 2x + 3x^2 - 4x^3 + \dots \\ &= 1 + (-1)2x + (-1)^2 3x^2 (-1)^3 4x^3 + \dots \end{aligned}$$

Following in this way we can write

$$= 1 + (-1)2x + (-1)^2 3x^2 (-1)^3 4x^3 + \dots + (-1)^n (n+1)x^n + \dots$$

So the term involving  $x^n = (-1)^n (n+1)x^n$

And hence coefficient of term involving  $x^n$  is  $(-1)^n (n+1)$

#### Question # 4

If  $x$  so small that its square and higher powers can be neglected, then show that

$$(i) \frac{1-x}{\sqrt{1+x}} \approx 1 - \frac{3}{2}x$$

$$(ii) \frac{\sqrt{1+2x}}{\sqrt{1-x}} \approx 1 + \frac{3}{2}x$$

$$(iii) \frac{(9+7x)^{\frac{1}{2}} - (16+3x)^{\frac{1}{4}}}{4+5x} \approx \frac{1}{4} - \frac{17}{384}x$$

$$(iv) \frac{\sqrt{4+x}}{(1-x)^3} \approx 2 + \frac{25}{4}x$$

$$(v) \frac{(1+x)^{\frac{1}{2}}(4-3x)^{\frac{1}{4}}}{(8+5x)^{\frac{1}{3}}} \approx \left(1 - \frac{5x}{6}\right)$$

$$(vi) \frac{(1-x)^{\frac{1}{2}}(9-4x)^{\frac{1}{2}}}{(8+3x)^{\frac{1}{3}}} \approx \frac{3}{2} - \frac{61}{48}x$$

$$(vii) \frac{\sqrt{4-x} + (8-x)^{\frac{1}{3}}}{(8-x)^{\frac{1}{3}}} \approx 2 - \frac{1}{12}x$$

**Solution**

(i)

$$\begin{aligned} \text{L.H.S} &= \frac{1-x}{\sqrt{1+x}} = \frac{1-x}{(1+x)^{\frac{1}{2}}} = (1-x)(1+x)^{-\frac{1}{2}} \\ &= (1-x) \left( 1 + \left( -\frac{1}{2} \right) (x) + \text{squares and higher power of } x \right). \\ &= 1 - x - \frac{1}{2}x + \text{squares and higher power of } x \\ &\approx 1 - \frac{3}{2}x = \text{R.H.S} \quad \text{Proved} \end{aligned}$$

$$(ii) \quad \text{Since} \quad \frac{\sqrt{1+2x}}{\sqrt{1-x}} = (1+2x)^{\frac{1}{2}}(1-x)^{-\frac{1}{2}}$$

$$\text{Now } (1+2x)^{\frac{1}{2}} = 1 + \left( \frac{1}{2} \right) (2x) + \text{squares and higher power of } x.$$

$$\begin{aligned} \text{Now } (1-x)^{-\frac{1}{2}} &= 1 + \left( -\frac{1}{2} \right) (-x) + \text{squares and higher power of } x. \\ &\approx 1 + \frac{1}{2}x \end{aligned}$$

$$\frac{\sqrt{1+2x}}{\sqrt{1-x}} \approx (1+x) \left( 1 + \frac{1}{2}x \right)$$

$$= 1 + x + \frac{1}{2}x \quad \text{ignoring term involving } x^2.$$

$$= 1 + \frac{3}{2}x \quad \text{Proved.}$$

$$(iii) \quad \frac{(9+7x)^{\frac{1}{2}} - (16+3x)^{\frac{1}{4}}}{4+5x} = \left( (9+7x)^{\frac{1}{2}} - (16+3x)^{\frac{1}{4}} \right) (4+5x)^{-1}$$

$$\text{Now } (9+7x)^{\frac{1}{2}} = 9^{\frac{1}{2}} \left( 1 + \frac{7x}{9} \right)^{\frac{1}{2}}$$

$$= (3^2)^{\frac{1}{2}} \left( 1 + \left( \frac{1}{2} \right) \left( \frac{7x}{9} \right) + \text{squares and higher of } x \right)$$

$$\begin{aligned}
 & \approx 3\left(1 + \frac{7x}{18}\right) = 3 + 3\left(\frac{7x}{18}\right) = 3 + \frac{7x}{6} \\
 (16+3x)^{\frac{1}{4}} &= (16)^{\frac{1}{4}}\left(1 + \frac{3x}{16}\right)^{\frac{1}{4}} \\
 &= (2^4)^{\frac{1}{4}}\left(1 + \left(\frac{1}{4}\right)\left(\frac{3x}{16}\right) + \text{square and higher power of } x\right) \\
 &\approx (2)\left(1 + \frac{3x}{64}\right) = 2 + 2\left(\frac{3x}{64}\right) = 2 + \frac{3x}{32} \\
 (4+5x)^{-1} &= 4^{-1}\left(1 + \frac{5}{4}x\right)^{-1} \\
 &= \frac{1}{4}\left(1 + (-1)\left(\frac{5}{4}x\right) + \text{squares and higher power of } x\right) \\
 &\approx \frac{1}{4}\left(1 - \frac{5}{4}x\right) = \frac{1}{4} - \frac{5}{16}x \\
 \text{So } \frac{(9+7x)^{\frac{1}{2}} - (16+3x)^{\frac{1}{4}}}{4+5x} &\approx \left[\left(3 + \frac{7x}{6}\right) - \left(2 + \frac{3x}{32}\right)\right]\left(\frac{1}{4} - \frac{5}{16}x\right) \\
 &= \left[3 + \frac{7x}{6} - 2 - \frac{3x}{32}\right]\left(\frac{1}{4} - \frac{5}{16}x\right) = \left(1 + \frac{103}{96}x\right)\left(\frac{1}{4} - \frac{5}{16}x\right) \\
 &= \frac{1}{4} + \frac{103}{384}x - \frac{5}{16}x = \frac{1}{4} - \frac{17}{384}x \quad \text{Proved}
 \end{aligned}$$

(iv) *Do yourself*

$$\begin{aligned}
 \text{(v)} \quad \frac{(1+x)^{\frac{1}{2}}(4-3x)^{\frac{3}{2}}}{(8+5x)^{\frac{1}{3}}} &= (1+x)^{\frac{1}{2}}(4-3x)^{\frac{3}{2}}(8+5x)^{-\frac{1}{3}} \\
 \text{Now } (1+x)^{\frac{1}{2}} &= 1 + \left(\frac{1}{2}\right)(x) + \text{square and higher power of } x \\
 &\approx 1 + \frac{1}{2}x \\
 (4-3x)^{\frac{3}{2}} &= 4^{\frac{3}{2}}\left(1 - \frac{3}{4}x\right)^{\frac{3}{2}} \\
 &= (2^2)^{\frac{3}{2}}\left(1 + \left(\frac{3}{2}\right)\left(-\frac{3}{4}x\right) + \text{square and higher power of } x\right) \\
 &\approx (2)^3\left(1 - \frac{9}{8}x\right) = 8\left(1 - \frac{9}{8}x\right) \\
 (8+5x)^{-\frac{1}{3}} &= (8)^{-\frac{1}{3}}\left(1 + \frac{5}{8}x\right)^{-\frac{1}{3}}
 \end{aligned}$$

$$\begin{aligned}
 &= (2^3)^{-\frac{1}{3}} \left( 1 + \left( -\frac{1}{3} \right) \left( \frac{5}{8}x \right) + \text{square and higher power of } x \right) \\
 &\approx (2)^{-1} \left( 1 - \frac{5}{24}x \right) = \frac{1}{2} \left( 1 - \frac{5}{24}x \right) \\
 \text{So } &\frac{(1+x)^{\frac{1}{2}}(4-3x)^{\frac{3}{2}}}{(8+5x)^{\frac{1}{3}}} \approx \left( 1 + \frac{1}{2}x \right) 8 \left( 1 - \frac{9}{8}x \right) \frac{1}{2} \left( 1 - \frac{5}{24}x \right) \\
 &= \frac{8}{2} \left( 1 + \frac{1}{2}x \right) \left( 1 - \frac{9}{8}x - \frac{5}{24}x \right) \\
 &= 4 \left( 1 + \frac{1}{2}x \right) \left( 1 - \frac{4}{3}x \right) = 4 \left( 1 + \frac{1}{2}x - \frac{4}{3}x \right) = 4 \left( 1 - \frac{5}{6}x \right) \text{ Proved}
 \end{aligned}$$

- (vi) *Do yourself as above*  
 (vii) *Same as Question #4 (iii)*

**Question # 5**

If  $x$  is so small that its cube and higher power can be neglected, then show that

$$(i) \sqrt{1-x-2x^2} = 1 - \frac{1}{2}x - \frac{9}{8}x^2 \quad (ii) \sqrt{\frac{1+x}{1-x}} = 1 + x + \frac{1}{2}x^2$$

**Solution**

$$\begin{aligned}
 (i) \sqrt{1-x-2x^2} &= (1-(x+2x^2))^{\frac{1}{2}} \\
 &= 1 + \left( \frac{1}{2} \right) (-x-2x^2) + \frac{\frac{1}{2}(\frac{1}{2}-1)}{2!} (-x-2x^2)^2 + \text{cube \& higher power of } x. \\
 &\approx 1 - \left( \frac{1}{2} \right) (x+2x^2) + \frac{\frac{1}{2}(-\frac{1}{2})}{2} (x+2x^2)^2 \\
 &\approx 1 - \frac{1}{2}x - \frac{1}{2}(2x^2) - \frac{1}{8}x^2 = 1 - \frac{1}{2}x - x^2 - \frac{1}{8}x^2 \\
 &= 1 - \frac{1}{2}x - \frac{9}{8}x^2 \quad \text{Proved}
 \end{aligned}$$

(ii)

$$\sqrt{\frac{1+x}{1-x}} = \frac{(1+x)^{\frac{1}{2}}}{(1-x)^{\frac{1}{2}}} = (1+x)^{\frac{1}{2}}(1-x)^{-\frac{1}{2}}$$

Now

$$\begin{aligned}
 (1+x)^{\frac{1}{2}} &= 1 + \left( \frac{1}{2} \right) x + \frac{\frac{1}{2}(\frac{1}{2}-1)}{2!} x^2 + \text{cube \& higher power of } x. \\
 &\approx 1 + \frac{1}{2}x + \frac{\frac{1}{2}(-\frac{1}{2})}{2} x^2 = 1 + \frac{1}{2}x - \frac{1}{8}x^2
 \end{aligned}$$

$$(1-x)^{-\frac{1}{2}} = 1 + \left(-\frac{1}{2}\right)(-x) + \frac{\left(-\frac{1}{2}\right)\left(-\frac{1}{2}-1\right)}{2!}(-x)^2 + \text{cube \& higher power of } x.$$

$$\approx 1 + \frac{1}{2}x + \frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)}{2}x^2 = 1 + \frac{1}{2}x + \frac{3}{8}x^2$$

So

$$\sqrt{\frac{1+x}{1-x}} = \left(1 + \frac{1}{2}x - \frac{1}{8}x^2\right) \left(1 + \frac{1}{2}x + \frac{3}{8}x^2\right)$$

$$= 1 + \frac{1}{2}x - \frac{1}{8}x^2 + \frac{1}{2}x + \frac{1}{4}x^2 + \frac{3}{8}x^2 = 1 + x + \frac{1}{2}x^2 \quad \text{Proved}$$

### Question # 6

If  $x$  is very nearly equal 1, then prove that  $px^p - qx^q = (p - q)x^{p+q}$

*Solution*

Since  $x$  is nearly equal to 1 so suppose  $x = 1 + h$ ,

where  $h$  is so small that its square and higher powers be neglected

$$\begin{aligned} \text{L.H.S} &= px^p - qx^q \\ &= p(1+h)^p - q(1+h)^q \\ &= p(1+ph + \text{square \& higher power of } x) \\ &\quad - q(1+qh + \text{square \& higher power of } h) \\ &\approx p(1+ph) - q(1+qh) \\ &= p + p^2h - q - q^2h \dots \text{(i)} \end{aligned}$$

$$\begin{aligned} \text{Now R.H.S} &= (p-q)x^{p+q} \\ &= (p-q)(1+h)^{p+q} \\ &= (p-q)(1+(p+q)h + \text{square \& higher power of } h) \\ &\approx (p-q)(1+(p+q)h) = (p-q)(1+ph+qh) \\ &= p + p^2h + pqh - q - pqh - q^2h \\ &= p + p^2h - q - q^2h \dots \text{(ii)} \end{aligned}$$

From (i) and (ii)

L.H.S  $\approx$  R.H.S Proved

### Question # 7

If  $p - q$  is small when compared with  $p$  or  $q$ , show that

$$\frac{(2n+1)p + (2n-1)q}{(2n-1)p + (2n+1)q} = \left(\frac{p+q}{2q}\right)^{\frac{1}{n}}.$$

*Solution* Since  $p - q$  is small when compare

Therefore let  $p - q = h \Rightarrow p = q + h$

$$\text{L.H.S} = \frac{(2n+1)p + (2n-1)q}{(2n-1)p + (2n+1)q} = \frac{(2n+1)(q+h) + (2n-1)q}{(2n-1)(q+h) + (2n+1)q}$$

$$\begin{aligned}
&= \frac{2nq + q + 2nh + h + 2nq - q}{2nq - q + 2nh - h + 2nq + q} = \frac{4nq + 2nh + h}{4nq + 2nh - h} \\
&= \frac{4nq + 2nh + h}{4nq \left(1 + \frac{2nh - h}{4nq}\right)} = \frac{4nq + 2nh + h}{4nq} \left(1 + \frac{2nh - h}{4nq}\right)^{-1} \\
&= \frac{4nq + 2nh + h}{4nq} \left(1 + (-1) \left(\frac{2nh - h}{4nq}\right) + \text{square \& higher power of } x^2\right) \\
&= \frac{4nq + 2nh + h}{4nq} \left(1 - \frac{2nh - h}{4nq}\right) = \frac{4nq + 2nh + h}{4nq} \left(\frac{4nq - 2nh + h}{4nq}\right) \\
&\approx \frac{16n^2q^2 + 8n^2hq + 4nhq - 8n^2hq + 4nhq}{16n^2q^2} \quad \text{ignoring squares of } h \\
&= \frac{16n^2q^2 + 8nhq}{16n^2q^2} = \frac{16n^2q^2}{16n^2q^2} + \frac{8nhq}{16n^2q^2} \\
&= 1 + \frac{h}{2nq} \dots \text{(i)}
\end{aligned}$$

$$\begin{aligned}
 \text{Now R.H.S} &= \left( \frac{p+q}{2q} \right)^{\frac{1}{n}} = \left( \frac{q+h+q}{2q} \right)^{\frac{1}{n}} \\
 &= \left( \frac{2q+h}{2q} \right)^{\frac{1}{n}} = \left( \frac{2q}{2q} + \frac{h}{2q} \right)^{\frac{1}{n}} = \left( 1 + \frac{h}{2q} \right)^{\frac{1}{n}} \\
 &= 1 + \left( \frac{1}{n} \right) \left( \frac{h}{2q} \right) + \text{square \& higher power of } h. \\
 &\approx 1 + \frac{h}{2nq} \quad \dots \dots \dots \text{ (ii)}
 \end{aligned}$$

Form (i) and (ii)

L.H.S  $\approx$  R.H.S      Proved

## Question # 8

Show that  $\left(\frac{n}{2(n+N)}\right)^{\frac{1}{2}} \approx \frac{8n}{9n-N} - \frac{n+N}{4n}$  where  $n$  and  $N$  are nearly equal.

**Solution** Since  $n$  and  $N$  are nearly equal therefore consider  $N = n + h$ , where  $h$  is so small that its squares and higher power be neglected.

$$\begin{aligned} \text{L.H.S.} &= \left( \frac{n}{2(n+N)} \right)^{\frac{1}{2}} = \left( \frac{n}{2(n+n+h)} \right)^{\frac{1}{2}} \\ &= \left( \frac{n}{2(2n+h)} \right)^{\frac{1}{2}} = \left( \frac{2(2n+h)}{n} \right)^{-\frac{1}{2}} = \left( \frac{4n+2h}{n} \right)^{-\frac{1}{2}} = \left( 4 + \frac{2h}{n} \right)^{-\frac{1}{2}} \end{aligned}$$

From (i) and (ii)

L.H.S = R.H.S      Proved

---

**Question # 9**

Identify the following series as binomial expansion and find the sum in each case.

$$(i) 1 - \frac{1}{2} \left( \frac{1}{4} \right) + \frac{1 \cdot 3}{2! \cdot 4} \left( \frac{1}{4} \right)^2 - \frac{1 \cdot 3 \cdot 5}{3! \cdot 8} \left( \frac{1}{4} \right)^3 + \dots$$

$$(ii) 1 - \frac{1}{2} \left( \frac{1}{2} \right) + \frac{1 \cdot 3}{2 \cdot 4} \left( \frac{1}{2} \right)^2 - \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} \left( \frac{1}{2} \right)^3 + \dots$$

$$(iii) \quad 1 + \frac{3}{4} + \frac{3 \cdot 5}{4 \cdot 8} + \frac{3 \cdot 5 \cdot 7}{4 \cdot 8 \cdot 12} + \dots$$

$$(iv) \quad 1 - \frac{1}{2} \left( \frac{1}{3} \right) + \frac{1 \cdot 3}{2 \cdot 4} \left( \frac{1}{3} \right)^2 - \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} \left( \frac{1}{3} \right)^3 + \dots$$

### **Solution**

$$(i) \quad 1 - \frac{1}{2} \left( \frac{1}{4} \right) + \frac{1 \cdot 3}{2! \cdot 4} \left( \frac{1}{4} \right)^2 - \frac{1 \cdot 3 \cdot 5}{3! \cdot 8} \left( \frac{1}{4} \right)^3 + \dots$$

Suppose the given series be identical with

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \dots$$

$$\text{This implies } nx = -\frac{1}{2} \left( \frac{1}{4} \right) \dots \dots \dots \quad (i)$$

$$\frac{n(n-1)}{2!}x^2 = \frac{1 \cdot 3}{2! \cdot 4} \left(\frac{1}{4}\right)^2 \dots \dots \dots \text{(ii)}$$

$$\text{From (i)} \quad nx = -\frac{1}{8} \Rightarrow x = -\frac{1}{8n} \dots \dots \dots \text{(iii)}$$

Putting value of  $x$  in (ii)

$$\begin{aligned} & \frac{n(n-1)}{2!} \left(-\frac{1}{8n}\right)^2 = \frac{1 \cdot 3}{2! \cdot 4} \left(\frac{1}{4}\right)^2 \\ & \Rightarrow \frac{n(n-1)}{2} \left(\frac{1}{64n^2}\right) = \frac{3}{2 \cdot 4} \left(\frac{1}{16}\right) \\ & \Rightarrow \frac{(n-1)}{128n} = \frac{3}{128} \Rightarrow (n-1) = \frac{3}{128} \cdot 128n \Rightarrow n-1 = 3n \\ & \Rightarrow n-3n=1 \Rightarrow -2n=1 \Rightarrow \boxed{n = -\frac{1}{2}} \end{aligned}$$

Putting value of  $n$  in equation (iii)

$$x = -\frac{1}{8\left(-\frac{1}{2}\right)} \Rightarrow \boxed{x = \frac{1}{4}}$$

So

$$(1+x)^n = \left(1 + \frac{1}{4}\right)^{-\frac{1}{2}} = \left(\frac{5}{4}\right)^{\frac{1}{2}} = \left(\frac{4}{5}\right)^{\frac{1}{2}} = \sqrt{\frac{4}{5}}$$

(ii) Do yourself as above

$$\text{(iii)} \quad 1 + \frac{3}{4} + \frac{3 \cdot 5}{4 \cdot 8} + \frac{3 \cdot 5 \cdot 7}{4 \cdot 8 \cdot 12} + \dots$$

Suppose the given series be identical with

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \dots$$

$$\text{This implies } nx = \frac{3}{4} \dots \dots \dots \text{(i)}$$

$$\frac{n(n-1)}{2!}x^2 = \frac{3 \cdot 5}{4 \cdot 8} \dots \dots \dots \text{(ii)}$$

$$\text{From (i)} \quad nx = \frac{3}{4} \Rightarrow x = \frac{3}{4n} \dots \dots \dots \text{(iii)}$$

Putting value of  $x$  in (ii)

$$\begin{aligned} & \frac{n(n-1)}{2!} \left(\frac{3}{4n}\right)^2 = \frac{3 \cdot 5}{4 \cdot 8} \Rightarrow \frac{n(n-1)}{2} \left(\frac{9}{16n^2}\right) = \frac{15}{32} \\ & \Rightarrow \frac{9(n-1)}{32n} = \frac{15}{32} \Rightarrow 9(n-1) = \frac{15}{32} \cdot 32n \Rightarrow 9n - 9 = 15n \end{aligned}$$

$$\Rightarrow 9n - 15n = 9 \Rightarrow -6n = 9 \Rightarrow n = -\frac{9}{6} \Rightarrow \boxed{n = -\frac{3}{2}}$$

Putting value of  $n$  in equation (iii)

$$x = -\frac{3}{4(-\frac{3}{2})} \Rightarrow \boxed{x = -\frac{1}{2}}$$

$$\text{So } (1+x)^n = \left(1 - \frac{1}{2}\right)^{-\frac{3}{2}} = \left(\frac{1}{2}\right)^{-\frac{3}{2}} = (2)^{\frac{3}{2}} = (\sqrt{2})^3 = 2\sqrt{2} \text{ Answer}$$

(iv) *Do yourself as above*

---

### Question # 10

Use binomial theorem to show that  $1 + \frac{1}{4} + \frac{1 \cdot 3}{4 \cdot 8} + \frac{1 \cdot 3 \cdot 5}{4 \cdot 8 \cdot 12} + \dots = \sqrt{2}$

$$\text{Solution} \quad 1 + \frac{1}{4} + \frac{1 \cdot 3}{4 \cdot 8} + \frac{1 \cdot 3 \cdot 5}{4 \cdot 8 \cdot 12} + \dots$$

Suppose the given series be identical with

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \dots$$

This implies

$$nx = \frac{1}{4} \dots \dots \dots \text{(i)}$$

$$\frac{n(n-1)}{2!}x^2 = \frac{1 \cdot 3}{4 \cdot 8} \dots \dots \dots \text{(ii)}$$

$$\text{From (i)} \quad nx = \frac{1}{4} \Rightarrow x = \frac{1}{4n} \dots \dots \dots \text{(iii)}$$

Putting value of  $x$  in (ii)

$$\begin{aligned} \frac{n(n-1)}{2!} \left(\frac{1}{4n}\right)^2 &= \frac{1 \cdot 3}{4 \cdot 8} \Rightarrow \frac{n(n-1)}{2} \left(\frac{1}{16n^2}\right) = \frac{3}{32} \\ \Rightarrow \frac{(n-1)}{32n} &= \frac{3}{32} \Rightarrow (n-1) = \frac{3}{32} \cdot 32n \Rightarrow n-1 = 3n \\ \Rightarrow n-3n &= 1 \Rightarrow -2n = 1 \Rightarrow \boxed{n = -\frac{1}{2}} \end{aligned}$$

Putting value of  $n$  in equation (iii)

$$x = \frac{1}{4(-\frac{1}{2})} \Rightarrow \boxed{x = -\frac{1}{2}}$$

$$\text{So } (1+x)^n = \left(1 - \frac{1}{2}\right)^{-\frac{1}{2}} = \left(\frac{1}{2}\right)^{-\frac{1}{2}} = (2)^{\frac{1}{2}} = \sqrt{2}$$

$$\text{Hence } 1 + \frac{1}{4} + \frac{1 \cdot 3}{4 \cdot 8} + \frac{1 \cdot 3 \cdot 5}{4 \cdot 8 \cdot 12} + \dots = \sqrt{2} \quad \text{Proved}$$


---

**Question # 11**

If  $y = \frac{1}{3} + \frac{1 \cdot 3}{2!} \left(\frac{1}{3}\right)^2 + \frac{1 \cdot 3 \cdot 5}{3!} \left(\frac{1}{3}\right)^3 + \dots$ , then prove that  $y^2 + 2y - 2 = 0$ .

$$\text{Solution} \quad y = \frac{1}{3} + \frac{1 \cdot 3}{2!} \left(\frac{1}{3}\right)^2 + \frac{1 \cdot 3 \cdot 5}{3!} \left(\frac{1}{3}\right)^3 + \dots$$

Adding 1 on both sides

$$1 + y = 1 + \frac{1}{3} + \frac{1 \cdot 3}{2!} \left(\frac{1}{3}\right)^2 + \frac{1 \cdot 3 \cdot 5}{3!} \left(\frac{1}{3}\right)^3 + \dots$$

Let the given series be identical with

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!} x^2 + \dots$$

This implies

$$nx = \frac{1}{3} \dots \dots \dots \quad (\text{i})$$

$$\frac{n(n-1)}{2!} x^2 = \frac{1 \cdot 3}{2!} \left(\frac{1}{3}\right)^2 \dots \dots \dots \quad (\text{ii})$$

$$\text{From (i)} \quad nx = \frac{1}{3} \Rightarrow x = \frac{1}{3n} \dots \dots \dots \quad (\text{iii})$$

Putting value of  $x$  in (ii)

$$\begin{aligned} \frac{n(n-1)}{2!} \left(\frac{1}{3n}\right)^2 &= \frac{1 \cdot 3}{2!} \left(\frac{1}{3}\right)^2 \\ \Rightarrow \frac{n(n-1)}{2} \left(\frac{1}{9n^2}\right) &= \frac{3}{2} \cdot \frac{1}{9} \\ \Rightarrow \frac{n-1}{18n} &= \frac{1}{6} \Rightarrow n-1 = \frac{1}{6} \cdot 18n \\ \Rightarrow n-1 &= 3n \Rightarrow n-3n = 1 \\ \Rightarrow -2n &= 1 \Rightarrow n = -\frac{1}{2} \end{aligned}$$

Putting value of  $n$  in equation (iii)

$$x = \frac{1}{3(-\frac{1}{2})} \Rightarrow x = -\frac{2}{3}$$

$$\begin{aligned} \text{So } (1+x)^n &= \left(1 - \frac{2}{3}\right)^{-\frac{1}{2}} = \left(\frac{1}{3}\right)^{-\frac{1}{2}} \\ &= (3)^{\frac{1}{2}} = \sqrt{3} \end{aligned}$$

This implies

$$1 + y = \sqrt{3}$$

On squaring both sides

$$\begin{aligned}(1+y)^2 &= (\sqrt{3})^2 \\ \Rightarrow 1+2y+y^2 &= 3 \Rightarrow 1+2y+y^2-3=0 \\ \Rightarrow y^2+2y-2 &= 0 \quad \text{Proved.}\end{aligned}$$


---

**Question # 12**

If  $2y = \frac{1}{2^2} + \frac{1 \cdot 3}{2!} \cdot \frac{1}{2^4} + \frac{1 \cdot 3 \cdot 5}{3!} \cdot \frac{1}{2^6} + \dots$ , then prove that  $4y^2 + 4y - 1 = 0$ .

**Solution**  $2y = \frac{1}{2^2} + \frac{1 \cdot 3}{2!} \cdot \frac{1}{2^4} + \frac{1 \cdot 3 \cdot 5}{3!} \cdot \frac{1}{2^6} + \dots$

Adding 1 on both sides

$$1+2y=1+\frac{1}{2^2}+\frac{1 \cdot 3}{2!} \cdot \frac{1}{2^4}+\frac{1 \cdot 3 \cdot 5}{3!} \cdot \frac{1}{2^6}+\dots$$

Let the given series be identical with

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \dots$$

This implies

$$nx = \frac{1}{2^2} \dots \dots \dots \text{(i)}$$

$$\frac{n(n-1)}{2!}x^2 = \frac{1 \cdot 3}{2!} \cdot \frac{1}{2^4} \dots \dots \text{(ii)}$$

From (i)  $nx = \frac{1}{4} \Rightarrow x = \frac{1}{4n} \dots \dots \text{(iii)}$

Putting value of  $x$  in (ii)

$$\begin{aligned}\frac{n(n-1)}{2!} \left( \frac{1}{4n} \right)^2 &= \frac{1 \cdot 3}{2!} \cdot \frac{1}{2^4} \\ \Rightarrow \frac{n(n-1)}{2} \left( \frac{1}{16n^2} \right) &= \frac{3}{2} \cdot \frac{1}{16} \\ \Rightarrow \frac{n-1}{n} &= 3 \Rightarrow n-1 = 3n\end{aligned}$$

$$\Rightarrow n-3n=1 \Rightarrow -2n=1 \Rightarrow \boxed{n=-\frac{1}{2}}$$

Putting value of  $n$  in equation (iii)

$$x = \frac{1}{4(-\frac{1}{2})} \Rightarrow \boxed{x = -\frac{1}{2}}$$

So

$$\begin{aligned}(1+x)^n &= \left(1-\frac{1}{2}\right)^{-\frac{1}{2}} \\ &= \left(\frac{1}{2}\right)^{-\frac{1}{2}} = (2)^{\frac{1}{2}} = \sqrt{2}\end{aligned}$$

This implies

$$1+2y=\sqrt{2}$$

On squaring both sides

$$\begin{aligned} (1+2y)^2 &= (\sqrt{2})^2 \\ \Rightarrow 1+4y+4y^2 &= 4 \quad \Rightarrow 1+4y+4y^2-2=0 \\ \Rightarrow 4y^2+4y-1 &= 0 \quad \text{Proved} \end{aligned}$$


---

### Question # 13

If  $y = \frac{2}{5} + \frac{1 \cdot 3}{2!} \left(\frac{2}{5}\right)^2 + \frac{1 \cdot 3 \cdot 5}{3!} \left(\frac{2}{5}\right)^3 + \dots$ , then prove that  $y^2 + 2y - 4 = 0$ .

**Solution**  $y = \frac{2}{5} + \frac{1 \cdot 3}{2!} \left(\frac{2}{5}\right)^2 + \frac{1 \cdot 3 \cdot 5}{3!} \left(\frac{2}{5}\right)^3 + \dots$

Adding 1 on both sides

$$1+y = 1 + \frac{2}{5} + \frac{1 \cdot 3}{2!} \left(\frac{2}{5}\right)^2 + \frac{1 \cdot 3 \cdot 5}{3!} \left(\frac{2}{5}\right)^3 + \dots$$

Let the given series be identical with

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!} x^2 + \dots$$

This implies

$$nx = \frac{2}{5} \dots \text{(i)}$$

$$\frac{n(n-1)}{2!} x^2 = \frac{1 \cdot 3}{2!} \left(\frac{2}{5}\right)^2 \dots \text{(ii)}$$

$$\text{From (i)} \quad nx = \frac{2}{5} \Rightarrow x = \frac{2}{5n} \dots \text{(iii)}$$

Putting value of  $x$  in (ii)

$$\begin{aligned} \frac{n(n-1)}{2!} \left(\frac{2}{5n}\right)^2 &= \frac{1 \cdot 3}{2!} \left(\frac{2}{5}\right)^2 \\ \Rightarrow \frac{n(n-1)}{2} \left(\frac{4}{25n^2}\right) &= \frac{3}{2} \left(\frac{4}{25}\right) \\ \Rightarrow \frac{n-1}{n} &= 3 \Rightarrow n-1 = 3n \Rightarrow n-3n = 1 \\ \Rightarrow -2n &= 1 \Rightarrow \boxed{n = -\frac{1}{2}} \end{aligned}$$

Putting value of  $n$  in equation (iii)

$$x = \frac{2}{5(-\frac{1}{2})} \Rightarrow \boxed{x = -\frac{4}{5}}$$

FSc-I / Ex 8.3 - 20

$$\text{So } (1+x)^n = \left(1 - \frac{4}{5}\right)^{-\frac{1}{2}} = \left(\frac{1}{5}\right)^{-\frac{1}{2}} = (5)^{\frac{1}{2}} = \sqrt{5}$$

This implies

$$1+y = \sqrt{5}$$

On squaring both sides

$$\begin{aligned} (1+y)^2 &= (\sqrt{5})^2 \\ \Rightarrow 1+2y+y^2 &= 5 \Rightarrow 1+2y+y^2-5 &= 0 \\ \Rightarrow y^2+2y-4 &= 0 \quad \text{Proved.} \end{aligned}$$

---